



PARISHRAMA NEET ACADEMY

☎ 080 4591 2222, Website: parishramaneetacademy.com

KCET – 2022 CHEMISTRY CODE B-3

PROVISIONAL ANSWER KEY - 2022

1. A first order reaction is half completed in 45 minutes. How long does it need 99.9% of the reaction to be completed?
 (A) 10 Hours (B) 20 Hours (C) 5 Hours **(D) 7.5 Hours**

Ans (D)

$$\begin{aligned} t_{99.9} &= 10t_{50} \\ &= 10 \times 45 \text{ min} \\ &= \frac{10 \times 45}{60} \text{ hr} \end{aligned}$$

2. The rate of the reaction: $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaOH} \rightarrow \text{CH}_3\text{COONa} + \text{C}_2\text{H}_5\text{OH}$ is given by the equation, $\text{Rate} = K [\text{CH}_3\text{COOC}_2\text{H}_5] [\text{NaOH}]$. If concentration is expressed in mol L^{-1} , the unit of K is
(A) $\text{L mol}^{-1} \text{s}^{-1}$ (B) s^{-1} (C) $\text{mol}^{-2} \text{L}^2 \text{s}^{-1}$ (D) $\text{mol L}^{-1} \text{s}^{-1}$

Ans (A)

$$\text{Rate} = k[\text{CH}_3\text{COOC}_2\text{H}_5]^1 [\text{NaOH}]^1$$

$$\text{Order} = 1 + 1 = 2$$

$$k_n = \text{mol}^{1-n} \cdot \text{L}^{n-1} \cdot \text{s}^{-1}$$

$$k_2 = \text{mol}^{-1} \cdot \text{L} \cdot \text{s}^{-1}$$

$$= \text{L} \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$$

3. Colloidal solution commonly used in the treatment of skin disease is
 (A) Colloidal Gold (B) Colloidal Antimony
(C) Colloidal Sulphur (D) Colloidal Silver

Ans (C)

Colloidal sulphur used in treatment of skin disease.



4. Specific conductance of 0.1 M HNO_3 is $6.3 \times 10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$. The molar conductance of the solution is

- (A) $6.300 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ (B) $63.0 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
(C) $630 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ (D) $315 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$

Ans (C)

$$\Lambda_m = \frac{\kappa \times 1000}{M}$$

$$= \frac{6.3 \times 10^{-2} \times 1000}{0.1} = 630 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

5. For spontaneity of a cell, which correct?

- (A) $\Delta G = +ve, \Delta E = +ve$ (B) **$\Delta G = -ve$**
 (C) $\Delta G = 0, \Delta E = 0$ (D) $\Delta G = -ve, \Delta E = 0$

Ans (B)

$$\Delta G = -nF(\Delta E)$$

For spontaneity of a cell $\Delta E = +ve > 0$

$$\Delta G = -ve < 0$$

6. For n^{th} order of reaction, Half-life period is directly proportional to

- (A) a^{n-1} (B) **a^{1-n}** (C) $\frac{1}{a^{n-1}}$ (D) $\frac{1}{a^{1-n}}$

Ans (B) and (C)

$$t_{1/2} \propto \frac{1}{a^{n-1}} \text{ or } t_{1/2} \propto a^{1-n}$$

7. Half-life of a reaction is found to be inversely proportional to the fifth power of its initial concentration, the order of reaction is

- (A) 5 (B) **6** (C) 3 (D) 4

Ans (B)

$$t_{1/2} \propto \frac{1}{a^{n-1}} \propto \frac{1}{a^5}$$

$$\therefore n - 1 = 5$$


$$n = 6$$

8. The strong reducing property of hypophosphorous acid is due to

- (A) two P-H bonds**
 (B) presence of phosphorus in its highest oxidation state

- (C) its concentration
(D) the positive valency of phosphorus

Ans (A)



Due to the presence of two P–H bonds.

9. A transition metal exists in its highest oxidation state. It is expected to behave as
(A) an oxidizing agent (B) a reducing agent
(C) a chelating agent (D) a central metal in a co-ordination compound

Ans (A)

For example KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$ used as oxidising agents.

10. What will be the value of x in Fe^{x+} , if the magnetic moment $\mu = \sqrt{24}$ BM?

- (A) 0 (B) +1 (C) +2 (D) +3

Ans (C)

Spin only magnetic moment = $\sqrt{n(n+2)}$ BM

Given magnetic moment = $\sqrt{24}$ Bm

$$\sqrt{n(n+2)} = \sqrt{24}$$

$$n = 4$$



11. Which can adsorb larger volume of hydrogen gas?

- (A) Finely divided platinum (B) Colloidal $\text{Fe}(\text{OH})_3$
(C) Finely divided nickel (D) Colloidal solution of palladium

Ans (D)

Surface area of palladium is maximum in its colloidal solution.

12. The property of halogens which is not correctly matched is

- (A) $\text{I} > \text{Br} > \text{Cl} > \text{F}$ (density)
(B) $\text{F} > \text{Cl} > \text{Br} > \text{I}$ (electron gain enthalpy)
(C) $\text{F} > \text{Cl} > \text{Br} > \text{I}$ (ionization enthalpy)
(D) $\text{F} > \text{Cl} > \text{Br} > \text{I}$ (electronegativity)

Ans (B)

$$\Delta H_{\text{eg}} = \text{Cl} > \text{F} > \text{Br} > \text{I}$$

F has lower electron gain enthalpy than chlorine due to small size 2p-orbital of fluorine.

13. Which noble gas has least tendency to form compounds?

- (A) Ar (B) Kr (C) He (D) Ne



Ans (C)

Due to small size and high ionization enthalpy of He.

14. $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ on heating liberates a gas. The same gas will be obtained by

- (A) treating H_2O_2 with NaNO_2 (B) treating Mg_3N_2 with H_2O
(C) heating NH_4NO_3 (D) heating NH_4NO_2


Ans (D)



15. The complex hexamine platinum (IV) chloride will give _____ number of ions on ionization.


- (A) 3 (B) 2 (C) 5 (D) 4

Ans (C)



Number of ions = one complex ion + four chloride ions

16. In the following pairs of halogen compounds, which compound undergoes faster $\text{S}_{\text{N}}1$ reaction?



Ans (D)

Reactivity order of alkyl halides in $\text{S}_{\text{N}}1$ reaction is $3^\circ > 2^\circ > 1^\circ$

17. The only Lanthanoid which is radioactive

- (A) Promethium (B) Praseodymium
(C) Lanthanum (D) Cerium

Ans (A)


Promethium is radioactive lanthanoid elements.

18. All Cu (II) halides are known, except the iodide, the reason for it is that

- (A) Cu^{2+} has much more negative hydration enthalpy.
 (B) Cu^{2+} ion has smaller size.
 (C) Iodide is bulky ion.

(D) Cu^{2+} oxidises iodide to iodine.

Ans (D)



19. The correct IUPAC name of *cis-platin* is

- (A) diammine dichlorido platinum (O)
 (B) dichlorido diammine platinum (IV)
(C) diammine dichlorido platinum (II)
 (D) diammine dichlorido platinum (IV)

Ans (C)

cis-platin is $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

IUPAC name: diammine dichloride platinum (II)

20. Crystal Field Splitting Energy (CFSE) for $[\text{CoCl}_6]^{4-}$ is 1800 cm^{-1} . The Crystal Field Splitting Energy (CFSE) for $[\text{CoCl}_6]^{4-}$ will be

- (A) 8000 cm^{-1}** (B) $10,000 \text{ cm}^{-1}$ (C) 18000 cm^{-1} (D) 16000 cm^{-1}


Ans (A)

$$\Delta t = \frac{4}{9} \Delta_o = \frac{4}{9} (18000) = 8,000 \text{ cm}^{-1}$$

21. The major product obtained when ethanol is heated with excess of concentrated H_2SO_4 at 443 K is

- (A) ethane (B) methane **(C) ethene** (D) ethyne


Ans (C)





22. Among the following, the products formed by the reaction of anisole with HI are

- (A) Benzene + Methanol (B) Phenol + Methane
(C) Phenol + Iodomethane (D) Sodium phenate + Methanol


Ans (C)



23. Which one of the following Chlorohydrocarbon readily undergoes solvolysis?



Ans (A)




Due to stable benzyl carbocation



24. Identify the products A and B in the reactions




Ans (A)



25. An organic compound with molecular formula $\text{C}_7\text{H}_8\text{O}$ dissolves in NaOH and gives a characteristic colour with FeCl_3 . On treatment with bromine, it gives a tribromo derivative $\text{C}_7\text{H}_5\text{OBr}_3$. The compound is



Ans (A)




Neutral FeCl_3 test is a characteristic test for phenolic group.

26. In Kolbes reaction the reacting substances are

- (A) Sodium phenate and CCl_4 (B) Phenol and CHCl_3
(C) Sodium phenate and CO_2 (D) Phenol and CCl_4


Ans (C)



27. In Carbylamine test for primary amines the resulting foul smelling product is

- (A) CH_3NC** (B) COCl_2 (C) CH_3NCl_2 (D) CH_3CN

Ans (A)



28. Ethanoic acid undergoes Hell-Volhard Zelinsky reaction but Methanoic acid does not, because of

- (A) absence of α -H atom in ethanoic acid
 (B) higher acidic strength of ethanoic acid than methanoic acid
 (C) presence of α -H atom in methanoic acid
(D) presence of α -H atom in ethanoic acid

Ans (D)

Condition to exhibit HVZ reaction is presence of α -hydrogen in a given carboxylic acid.

29. The general name of the compound formed the reaction between aldehyde and alcohol is

- (A) Glycol (B) Acetate (C) Ester **(D) Acetal**

Ans (D)

Nucleophilic condition of aldehydes and ketones forms corresponding hemiacetals and acetals.

30. Reaction by which benzaldehyde cannot be prepared is

- (A) Toluene $\xrightarrow[\text{(ii) H}_3\text{O}^+]{\text{(i) CrO}_2\text{Cl}_2 \text{ in CS}_2}$
 (B) Benzoyl chloride + $\text{H}_2 \xrightarrow[\text{heat}]{\text{Pd-BaSO}_4}$
 (C) Benzene + $\text{CO} + \text{HCl} \xrightarrow{\text{anhydrous AlCl}_3}$
(D) Benzoic acid $\xrightarrow{\text{Zn-Hg and conc. HCl}}$

Ans (D)

Option (A), (B) and (C) are the methods of preparation of benzaldehyde.

31. The test to differentiate between pentan-2-one and pentan-3-one is

- (A) Fehling's test (B) Iodoform test
(C) Baeyer's test (D) Benedict's tests

Ans (B)


Iodoform test is useful to differentiate pentan-2-one and pentan-3-one because pentan-2-one has methyl keto group but not pentan-3-one.

32. A secondary amine is

- (A) a compound with an NH_2 group on the carbon atom in number 2 position
(B) a compound in which 2 of the hydrogen of NH_3 have been replaced by organic groups
(C) an organic compound with two NH_2 group
(D) a compound with two carbon atom and an NH_2 group

Ans (B)

Amines are classified based on attachment of number of alkyl or aryl groups attached to the N.



33. Which of the following is correctly matched?

- (A) Bakelite – Novolac (B) Polyester – tetrafluoroethene
(C) Nylon – acrylonitrile (D) Teflon – copralactum

Ans (A)

Novolac on further polymerization produces bakelite.

34. Which institute has approved the emergency use of 2-deoxy-D-Glucose as additive therapy for COVID-19 patients?

- (A) Ministry of Health and Family Welfare
(B) Drug Controller General of India
(C) Indian Council of Medical Research
(D) World Health Organisation

Ans (B)

Drug controller general of India has approved the emergency use of 2-deoxy-D-Glucose.

35. A Nucleic acid, whether DNA or RNA gives on complete hydrolysis, two purine bases, two pyrimidine bases, a pentose sugar and phosphoric. Nucleotides which are intermediate products in the hydrolysis contain

- (A) purine or pyrimidine base and ortho-phosphoric acid
(B) Purine or pyrimidine base, a pentose sugar and ortho-phosphoric acid
 © purine or pyrimidine base and pentose sugar.
 (D) a purine base, pentose sugar and ortho-phosphoric acid

Ans (B)

Nucleic acid on hydrolysis produces nitrogenous base + pentose sugar + ortho phosphoric acid.

36. Which is most VISCOUS?

- (A) Ethylene glycol **(B) Glycerol**
 (C) Methanol (D) Ethanol

Ans (B)

Viscosity \propto force of attraction (H-bonding)

37. The volume of 2.8 g of CO at 27 °C and 0.821 atm pressure is

(R = 0.08210 L atm K⁻¹ mol⁻¹)

- (A) 3 litres** (B) 30 litres (C) 0.3 litres (D) 1.5 litres

Ans (A)

$$V = \frac{nRT}{P}$$

$$n_{\text{CO}} = \frac{2.8}{28} = 0.1$$

$$V = \frac{0.1 \times 0.0821 \times 300}{0.821} = 3 \text{ litres}$$

38. The work done when 2 moles of an ideal gas expands reversibly and isothermally from a volume of 1 L to 10 L at 300 K is (R = 0.0083 kJ K mol⁻¹)

- (A) 0.115 kJ (B) 58.5 kJ **(C) 11.5 kJ** (D) 5.8 kJ

Ans (C)

$$W = 2.303nRT \log_{10} \left[\frac{V_2}{V_1} \right]$$

$$= 2.303 \times 2 \times 0.0083 \times 300 \times \log_{10} \left(\frac{10}{1} \right) = 11.5 \text{ kJ}$$

39. An aqueous solution of alcohol contains 18 g of water and 414 g of alcohol. The mole fraction water is

- (A) 0.7 (B) 0.9 (C) 0.1 (D) 0.4

Ans (C)

$$\chi_{\text{H}_2\text{O}} = \frac{n_{\text{H}_2\text{O}}}{n_{\text{H}_2\text{O}} + n_{\text{C}_2\text{H}_5\text{OH}}} \quad n_{\text{H}_2\text{O}} = \frac{18}{18} = 1$$

$$= \frac{1}{1+9} = 0.1 \quad n_{\text{C}_2\text{H}_5\text{OH}} = \frac{414}{46} = 9$$

40. If wavelength of photon is 2.2×10^{-11} m and $h = 6.6 \times 10^{-34}$ s, then momentum of photon

- (A) 1.452×10^{-44} kg ms⁻¹ (B) $6.89 \times 10^{+43}$ kg ms⁻¹

- (C) 3×10^{-23} kg ms⁻¹ (D) 3.33×10^{-22} kg ms⁻¹

Ans (C)

$$\lambda = \frac{h}{p} \Rightarrow p = \frac{h}{\lambda}$$

$$p = \frac{6.6 \times 10^{-34}}{2.2 \times 10^{-11}} = 3 \times 10^{-23}$$

41. Elements X, Y and Z have atomic numbers 19, 37 and 35 respectively. Which of the following statements is true about them?

- (A) Z would have the highest ionization potential.
 (B) Y would have the highest ionization potential.
 (C) Their ionization potential, would increase with increasing atomic number.

(D) Y would have an ionization potential between those of X and Z.

Ans (D)


Down to the group of ionization enthalpy decreases.

42. In oxygen and carbon molecule the bonding is

- (A) O₂ : 1σ, 1π; C₂ : 0σ, 2π (B) O₂ : 0σ, 2π; C₂ : 2σ, 0π

- (C) O₂ : 1σ, 1π; C₂ : 1σ, 1π (D) O₂ : 2σ, 0π; C₂ : 0σ, 2π

Ans (A)



43. Amphoteric oxide among the following:

- (A) Ag₂O (B) SnO₂ (C) BeO (D) CO₂

Ans (B) and (C)

BeO-Amphoteric oxide.

44. Which property of CO₂ makes it biologically and geo-chemically important?

- (A) Its low solubility in water (B) Its high compressibility
(C) Its acidic nature (D) Its colourless and odourless nature

Ans (A)

Its low solubility in water

Due to low solubility of CO₂ in water, it is important in biological and geochemical process.

45. The IUPAC name for $\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_2 - \text{CH}_2 - \overset{\text{O}}{\parallel}{\text{C}} - \text{O} - \text{H}$ is

- (A) 1-carboxybutan-3-one (B) **4-oxopentanoic acid**
(C) 1-hydroxy pentane-1, 4-dione (D) 1, 4-dioxopentanol


Ans (B)

4-oxopentanoic acid.

46. 1 mole of HI is heated in a closed container of capacity of 2 L. At equilibrium half a mole of HI is dissociated. The equilibrium constant of the reaction is

- (A) 0.25 (B) 0.35 (C) **1** (D) 0.5

Ans (C)



Initial 1 0 0

At equ. ½ ¼ ¼

$$k = \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2}$$

$$k = \frac{\left(\frac{1}{4}\right)\left(\frac{1}{4}\right)}{\left(\frac{1}{2}\right)^2} = \frac{1}{4} = 0.25$$

47. Which among the following has highest pH?

- (A) 1 M H₂SO₄ (B) 0.1 M NaOH
(C) 1 M HCl (D) **1 M NaOH**

Ans (D)


1 M NaOH; [OH⁻] 1, pOH = -log 1 = 0

∴ pH = 14

48. In which of the following compounds, an element exhibits two different oxidation states?

- (A) N_2H_4 (B) N_3H (C) NH_2CONH_2 (D) NH_4NO_3

Ans (D)




49. Which of the following hydrides is electron deficient?

- (A) CH_4 (B) B_2H_6 (C) NaH (D) CaH_2


Ans (B)

B_2H_6 [13th group hydrides are electron deficient and also Lewi's acids]

50. Identify A and B in the reaction



Ans (D)



51. Vacant space in body centered cubic lattice unit cell is about

- (A) 23% (B) 46% (C) 32% (D) 10%

Ans (C)

BCC-void space – 32%

52. How many number of atoms are there in a cube based unit cell having one atom on each corner and 2 atom on each body diagonal of cube?

- (A) 4 (B) 9 (C) 8 (D) 6

Ans (B)

$$\text{Each corner} = 8 \times \frac{1}{8} = 1$$

$$4 \text{ body diagonals} = 4 \times 2 = 8$$

$$\text{Total} = 9$$

53. Which of the following is NOT true about the amorphous solids?

- (A) Amorphous solids can be moulded by heating.
(B) They are anisotropic in nature.
(C) On heating they may become crystalline at certain temperature.
(D) They may become crystalline on keeping for long time.

Ans (B)

They are anisotropic in nature.

54. Which of the following colligative properties can provide molar mass of proteins, polymers, and colloids with greater precision?

- (A) Depression in freezing point
(B) Osmotic pressure
(C) Relative lowering of vapour pressure
(D) Elevation in boiling point

Ans (B)

Osmotic pressure can provide molar mass of proteins, polymers and colloids.

55. In Fuel cells _____ are used as catalysts

- (A) Zinc - Mercury (B) Lead - Manganese
(C) Platinum - Palladium (D) Nickel - Cadmium

Ans (C)

Pt, Pd used as catalysis in fuel cells.

56. The molar conductivity is maximum for the solution of concentration

- (A) 0.005 M (B) 0.001 M (C) 0.004 M (D) 0.002 M

Ans (B)

$$\Lambda_m = \frac{\kappa \times 1000}{M}$$

$$\Lambda_m \propto \frac{1}{\text{Molarity}} \propto \text{Volume}$$

57. Alkali halides do not show dislocation defect because

(A) Cations and anions have almost equal size.

(B) There is large difference in size of cations and anions.

(C) Cations and anions have low co-ordination number.

(D) Anions cannot be accommodated in vacant spaces.

Ans (A)

58. Solubility of a gas in a liquid increases with

(A) increase of P and decrease of T (B) decrease of P and decrease of T

(C) increase of P and increase of T (D) decrease of P and increase of T

Ans (A)

Increase of P and decrease of T.

59. The rise in boiling point of a solution containing 1.8 g of glucose in 100 g of solvent is 1 °C. The molal elevation constant of the liquid is

(A) 2 K kg/mol (B) 10 K kg/mol (C) 0.1 K kg/mol (D) 1 K kg/mol

Ans (D)

$$\Delta T_b = i \times k_b \times m$$

$$0.1 = 1 \times k_b \times \frac{1.8}{180} \times \frac{1000}{100} \Rightarrow k_b = 1$$

60. If 3 g of glucose (molar mass = 180 g) is dissolved in 60 g of water at 15 °C, the osmotic pressure of the solution will be

(A) 6.57 atm (B) 5.57 atm (C) 0.34 atm (D) 0.65 atm

Ans (A)

$$\pi = i \times c \times R \times T = 1 \times \frac{3}{180} \times \frac{1000}{60} \times 0.0821 \times 288 = 6.57 \text{ atm}$$

* * *

PARISHRAMA
NEET ACADEMY



ಪರಿಶ್ರಮ
ನಿಜ್ಞಾ ಆಕಾಡೆಮಿ

PARISHRAMA NEET ACADEMY

Omkar Hills Road, Srinivasapura, Dr. Vishnuvardan Road, towards,
Uttarahalli Main Road, Bengaluru, 560060

☎ 080 4591 2222, Website: parishramaneetacademy.com

**ADMISSIONS OPEN
FOR 2022-23**

**FOR LONG TERM
NEET COURSE**

**CLASSES START FROM
18-JULY-2022**

NEET 220 Medical Seats out of 240

NEET 220 Medical Seats out of 240

FOR ENQUIRIES CONTACT

96866 12464

89042 72143

96115 60724

96115 60640

96115 60629

89047 67771

